

GEOMETRIC HEIGHT INEQUALITY ON VARIETIES WITH AMPLE COTANGENT BUNDLES

ATSUSHI MORIWAKI

November, 1993 (revised)

ABSTRACT. Let F be a function field of one variable over an algebraically closed field of characteristic zero, X a geometrically irreducible smooth projective variety over F , and L a line bundle on X . In this note, we will prove that if $\Omega_{X/F}^1$ is ample and X is non-isotrivial, then there are a proper closed algebraic set Y of X and a constant $A > 0$ such that

$$h_L(P) \leq A \cdot d(P) + O(1)$$

for all $P \in X(\bar{F}) \setminus Y(\bar{F})$, where $h_L(P)$ is a geometric height of P with respect to L and $d(P)$ is the geometric logarithmic discriminant of P . As corollary of the above height inequality, we can recover Noguchi's theorem [No], i.e. there is a non-empty Zariski open set U of X with $U(F) = \emptyset$.

0. Introduction.

Let k be an algebraically closed field of characteristic zero and F a finitely generated extension field of k with $\text{trans. deg}_k F = 1$. We will fix two fields k and F throughout this note.

Let X be a geometrically irreducible smooth projective variety over F . Let $\mathcal{X}$ and C be smooth projective varieties over k , and $f : \mathcal{X} \rightarrow C$ a k -morphism such that the function field of C is F and the generic fiber of f is X , i.e. $X = \mathcal{X} \otimes F$. In this note, X is said to be *non-isotrivial* if there is a non-empty open set C_0 of C such that, for all $t \in C_0$, the Kodaira-Spencer map $\varrho_t : T_{C,t} \rightarrow H^1(\mathcal{X}_t, T_{\mathcal{X}_t})$ is not zero. Let $\bar{F}$ be the algebraic closure of F . For a point $P \in X(\bar{F})$, let us denote by Δ_P the corresponding integral curve on $\mathcal{X}$. We fix a line bundle L on X . Let $\mathcal{L}$ be a line bundle on $\mathcal{X}$ with $\mathcal{L} \otimes F = L$. The pair $(f : \mathcal{X} \rightarrow C, \mathcal{L})$ is called a *model of (X, L)* . A *geometric height $h_L(P)$ of P with respect to L* is defined by

$$h_L(P) = \frac{(\mathcal{L} \cdot \Delta_P)}{[F(P) : F]}.$$

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$

The geometric height h_L might depend on the choice of the model $(f : \mathcal{X} \rightarrow C, \mathcal{L})$ of (X, L) , but as we will see in Proposition 1.1, it is uniquely determined modulo bounded functions. Moreover, the geometric logarithmic discriminant $d(P)$ of P is given by

$$d(P) = \frac{2g(\tilde{\Delta}_P) - 2}{[F(P) : F]},$$

where $g(\tilde{\Delta}_P)$ is the genus of the normalization $\tilde{\Delta}_P$ of Δ_P . An interesting problem concerning h_L and d on the non-isotrivial X is to find an inequality of type:

$$h_L(P) \leq A \cdot d(P) + O(1)$$

for all $P \in X(\bar{F})$, which is called a *geometric height inequality*. In the case where $\dim X = 1$, we know a lot of height inequalities as follows, where $g = g(X) \geq 2$ and $K = K_{\mathcal{X}/C}$.

L. Szpiro,	$h_K(P) \leq 24(g-1)^2 d(P) + O(1),$	(char ≥ 0)
P. Vojta,	$h_K(P) \leq (8g-6)/3 d(P) + O(1),$	(char $= 0$)
A. N. Parshin,	$h_K(P) \leq (20g-15)/6 d(P) + O(1),$	(char $= 0$)
H. Esnault and E. Viehweg,	$h_K(P) \leq 2(2g-1)^2 d(P) + O(1),$	(char $= 0$)
P. Vojta,	$h_K(P) \leq (2+\epsilon) d(P) + O(1),$	(char $= 0$)
A. Moriwaki,	$h_K(P) \leq (2g-1) d(P) + O(1),$	(char ≥ 0)

Details can be found in Chapter VI of [La] or [Mo]. In this note, inspired by [Vo], I would like to study a geometric height inequality on a higher dimensional variety. The main theorem of this note is

Theorem A. *Let X be a geometrically irreducible smooth projective variety over F and L a line bundle on X . If $\Omega_{X/F}^1$ is ample and X is non-isotrivial, then there are a proper closed algebraic set Y of X and a constant $A > 0$ such that*

$$h_L(P) \leq A \cdot d(P) + O(1)$$

for all $P \in X(\bar{F}) \setminus Y(\bar{F})$.

Under the assumption of Theorem A, it might be not needed to subtract the proper closed algebraic set Y . But, it is sufficient to recover Noguchi's theorem [No] concerning higher dimensional geometric Mordell's conjecture, i.e.

Theorem B. *Let X be a geometrically irreducible smooth projective variety over F . If $\Omega_{X/F}^1$ is ample and X is non-isotrivial, then there is a non-empty Zariski open set U of X with $U(F) = \emptyset$.*

Theorem B is a corollary of Theorem A once we get finiteness property of geometric heights. However, finiteness does not hold in general for geometric heights. Instead of finiteness, algebraic heights have non-denseness property (cf. Theorem 1.3).

In §1, we will deal with properties of geometric heights, especially 'positivity' and 'non-denseness'. In §2, we will prepare several basic facts of ample vector bundles. §3 is devoted to the proofs of Theorem A and Theorem B.

1. Properties of geometric heights.

Let X be a geometrically irreducible smooth projective variety over F and L a line bundle on X . First of all, we must check the following proposition. By this proposition,

$$h_L(P) = \frac{(\mathcal{L} \cdot \Delta_P)}{[F(P) : F]}$$

is uniquely determined modulo bounded functions.

Proposition 1.1. *Let $(f : \mathcal{X} \rightarrow C, \mathcal{L})$ and $(f' : \mathcal{X}' \rightarrow C, \mathcal{L}')$ be two models of (X, L) . Then, there is a constant A such that, for all $P \in X(\bar{F})$,*

$$\left| \frac{(\mathcal{L} \cdot \Delta_P)}{[F(P) : F]} - \frac{(\mathcal{L}' \cdot \Delta'_P)}{[F(P) : F]} \right| \leq A.$$

Proof. We can easily construct birational morphisms $\mu : \mathcal{X}'' \rightarrow \mathcal{X}$ and $\mu' : \mathcal{X}'' \rightarrow \mathcal{X}'$ such that $(\mathcal{X}'' \rightarrow C, \mu^*(\mathcal{L}))$ and $(\mathcal{X}'' \rightarrow C, \mu'^*(\mathcal{L}'))$ are models of (X, L) . So we may assume that $\mathcal{X} = \mathcal{X}'$.

Since $\mathcal{L} \otimes \mathcal{L}'^{-1}$ is trivial on X , we have $f_*(\mathcal{L} \otimes \mathcal{L}'^{-1}) \neq 0$. Thus there is an ample line bundle A_1 on C such that $H^0(\mathcal{X}, \mathcal{L} \otimes \mathcal{L}'^{-1} \otimes f^*(A_1)) \neq 0$. Let D be an element of $|\mathcal{L} \otimes \mathcal{L}'^{-1} \otimes f^*(A_1)|$. D has no horizontal components because $D|_X$ is linearly equivalent to zero. Thus, for all $P \in X(\bar{F})$, $(D \cdot \Delta_P) \geq 0$. It follows that

$$\frac{(\mathcal{L}' \cdot \Delta_P)}{[F(P) : F]} - \frac{(\mathcal{L} \cdot \Delta_P)}{[F(P) : F]} \leq \deg(A_1).$$

By the same way, we can find an ample line bundle A_2 on C such that

$$\frac{(\mathcal{L} \cdot \Delta_P)}{[F(P) : F]} - \frac{(\mathcal{L}' \cdot \Delta_P)}{[F(P) : F]} \leq \deg(A_2).$$

Hence, we get our proposition. $\square$

Let V be a projective variety defined over an algebraic number field K , L a line bundle on V , and h_L a Weil-height of L . Then, h_L has the following two basic properties:

- (1) (Positivity) Let B be a base locus of $|L|$. Then, $h_L(P) \geq O(1)$ for all $P \in V(\bar{\mathbb{Q}}) \setminus B(\bar{\mathbb{Q}})$.
- (2) (Finiteness) If L is ample and $A \in \mathbb{R}$, then the set $\{P \in V(K) \mid h_L(P) \leq A\}$ is finite.

The main topic of this section is to consider geometric analogies of the above two properties of arithmetic heights. First, let us consider positivity of geometric heights.

Proposition 1.2. (*Positivity*) *Let X be a geometrically irreducible smooth projective variety over F and L a line bundle on X . Then, $h_L(P) \geq O(1)$ for all $P \in X(\bar{F}) \setminus \text{Bs}(L)(\bar{F})$, where $\text{Bs}(L) = \text{Supp}(\text{Coker}(H^0(X, L) \otimes \mathcal{O}_X \rightarrow L))$.*

Proof. Let $(f : \mathcal{X} \rightarrow C, \mathcal{L})$ be a model of (X, L) . Let A be an ample line bundle on C such that $f_*(\mathcal{L}) \otimes A$ is generated by global sections. Let $P \in X(\bar{F}) \setminus \text{Bs}(L)(\bar{F})$ and Δ_P the corresponding integral curve on $\mathcal{X}$. Assume that $\Delta_P \subset \text{Bs}(\mathcal{L} \otimes f^*(A))$. Then, the natural homomorphism

$$H^0(C, f_*(\mathcal{L} \otimes f^*(A) \otimes I_{\Delta_P})) \rightarrow H^0(C, f_*(\mathcal{L} \otimes f^*(A)))$$

is surjective, where I_{Δ_P} is the defining ideal of Δ_P . Here, we consider the following commutative diagram:

$$\begin{array}{ccc} H^0(C, f_*(\mathcal{L} \otimes f^*(A) \otimes I_{\Delta_P})) \otimes \mathcal{O}_C & \xrightarrow{\alpha} & H^0(C, f_*(\mathcal{L} \otimes f^*(A))) \otimes \mathcal{O}_C \\ \gamma \downarrow & & \downarrow \delta \\ f_*(\mathcal{L} \otimes f^*(A) \otimes I_{\Delta_P}) & \xrightarrow[\beta]{} & f_*(\mathcal{L} \otimes f^*(A)) \end{array}$$

Since α and δ are surjective, so is β , which implies that $H^0(Y_{\bar{F}}, L_{\bar{F}} \otimes I_P) \rightarrow H^0(Y_{\bar{F}}, L_{\bar{F}})$ is surjective. Therefore, $P \in \text{Bs}(L_{\bar{F}})$. This is a contradiction. Thus, we get $\Delta_P \not\subset \text{Bs}(\mathcal{L} \otimes f^*(A))$. Hence, $(\mathcal{L} \otimes f^*(A) \cdot \Delta_P) \geq 0$, which says that $h_L(P) \geq -\deg A$. $\square$

Next, we will consider finiteness of geometric heights. Unfortunately, finiteness does not hold in general for geometric heights. Instead of it, we have the following non-denseness of geometric heights.

Theorem 1.3. (*Non-denseness*) *Let X be a geometrically irreducible smooth projective variety over F and L an ample line bundle on X . We assume that X contains no rational curves and X is non-isotrivial. Then, for every constant A , the set $\{P \in X(F) \mid h_L(P) \leq A\}$ is not dense in $X(\bar{F})$.*

First of all, we will prepare two basic lemmas.

Lemma 1.4. *Let $q : Y \rightarrow X$ be a surjective morphism of normal projective varieties over an algebraically closed field of characteristic zero and E a vector bundle on X . Then, the natural morphism $H^1(X, E) \rightarrow H^1(Y, q^*E)$ is injective.*

Proof. Let us consider the Leray spectral sequence:

$$E_2^{i,j} = H^i(X, R^j q_*(q^*E)) \implies E^{i+j} = H^{i+j}(Y, q^*E).$$

Since

$$E_2^{1,0} = H^1(X, q_*(q^*E)) \rightarrow E^1 = H^1(Y, q^*E)$$

is injective, it is sufficient to see that

$$H^1(X, E) \rightarrow H^1(X, q_*(\mathcal{O}_Y) \otimes E)$$

is injective. Let

$$Y \xrightarrow{q_1} Z \xrightarrow{q_2} X$$

be the Stein factorization of $q : Y \rightarrow X$. Then, $q_*(\mathcal{O}_Y) = q_{2*}(\mathcal{O}_Z)$. Thus, we may assume that q is a finite morphism. Since we work over the field of characteristic zero, we have a trace map

$$\mathrm{Tr} : q_*\mathcal{O}_Y \rightarrow \mathcal{O}_X$$

such that $\mathrm{Tr} \cdot \iota = \mathrm{id}$, where ι is the inclusion map $\iota : \mathcal{O}_X \rightarrow q_*(\mathcal{O}_Y)$. This means that the exact sequence

$$0 \rightarrow E \rightarrow q_*(\mathcal{O}_Y) \otimes E \rightarrow (q_*(\mathcal{O}_Y)/\mathcal{O}_X) \otimes E \rightarrow 0$$

splits. Therefore, $H^1(X, E) \rightarrow H^1(X, q_*(\mathcal{O}_Y) \otimes E)$ is injective. $\square$

Lemma 1.5. *Let $\mathcal{X}$ be a smooth projective variety over k , C a smooth projective curve over k , $f : \mathcal{X} \rightarrow C$ a k -morphism with $f_*\mathcal{O}_{\mathcal{X}} = \mathcal{O}_C$. Assume that there are a non-empty open set C_0 of C , a smooth projective variety Y over k and a dominant morphism $\phi : Y \times C_0 \rightarrow \mathcal{X}_0 = f^{-1}(C_0)$ such that $p = f_0 \cdot \phi$, where $p : Y \times C_0 \rightarrow C_0$ is the natural projection and $f_0 = f|_{\mathcal{X}_0}$.*

$$\begin{array}{ccc} Y \times C_0 & \xrightarrow{\phi} & \mathcal{X}_0 \\ p \downarrow & & \downarrow f_0 \\ C_0 & \xlongequal{\quad} & C_0 \end{array}$$

Then, for a general point t of C_0 , the Kodaira-Spencer map $\varrho_t : T_{C,t} \rightarrow H^1(\mathcal{X}_t, T_{\mathcal{X}_t})$ is zero.

Proof. Shrinking C_0 , if necessarily, we may assume that f_0 is smooth. Let us consider the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & T_{Y \times C_0 / C_0} & \longrightarrow & T_{Y \times C_0} & \longrightarrow & p^*(T_{C_0}) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \phi^*(T_{\mathcal{X}_0 / C_0}) & \longrightarrow & \phi^*(T_{\mathcal{X}_0}) & \longrightarrow & \phi^*(f_0^*(T_{C_0})) \longrightarrow 0 \end{array}$$

This gives rise to a commutative diagram:

$$\begin{array}{ccc} T_{C,t} & \xrightarrow{\alpha_t} & H^1(Y \times \{t\}, T_{Y \times \{t\}}) \\ \parallel & & \downarrow \\ T_{C,t} & \xrightarrow{\beta_t} & H^1(Y \times \{t\}, \phi_t^*(T_{\mathcal{X}_t})) \end{array}$$

for a point $t \in C_0$. Since $\alpha_t = 0$, we have $\beta_t = 0$. Here β_t is factored through the Kodaira-Spencer map $\varrho_t : T_{C,t} \rightarrow H^1(\mathcal{X}_t, T_{\mathcal{X}_t})$, i.e. if we denote by γ_t the natural homomorphism $H^1(\mathcal{X}_t, T_{\mathcal{X}_t}) \rightarrow H^1(Y \times \{t\}, \phi_t^*(T_{\mathcal{X}_t}))$, then $\beta_t = \gamma_t \cdot \varrho_t$.

$$T_{C,t} \xrightarrow{\varrho_t} H^1(\mathcal{X}_t, T_{\mathcal{X}_t}) \xrightarrow{\gamma_t} H^1(Y \times \{t\}, \phi_t^*(T_{\mathcal{X}_t}))$$

On the other hand, by Lemma 1.4, γ_t is injective. Thus, $\varrho_t = 0$ $\square$

Proof of Theorem 1.3. Let $(f : \mathcal{X} \rightarrow C, \mathcal{L})$ be a model of (X, L) . Let $\mathcal{M}$ be an ample line bundle on $\mathcal{X}$. Since L is ample on X , for a sufficiently large n , $\mathcal{L}^n \otimes \mathcal{M}^{-1}$ is very ample on X . Thus, by Proposition 1.1, there is a constant B such that $h_{\mathcal{L}^n \otimes \mathcal{M}^{-1}}(P) \geq B$ for all $P \in X(\bar{F})$, where $M = \mathcal{M}_F$. Therefore, we have

$$h_M(P) \leq n \cdot h_L(P) - B$$

for all $P \in X(\bar{F})$. It follows that we may assume that $\mathcal{L}$ is ample on $\mathcal{X}$.

We set $S_A = \{P \in X(F) \mid h_L(P) \leq A\}$ and $\Sigma_A = \{\Delta_P \subset \mathcal{X} \mid P \in S_A\}$. Here we assume that S_A is dense in $X(\bar{F})$. First of all, we claim that there is a smooth quasi-projective variety Y over k and a dominant morphism $\phi : Y \times C \rightarrow \mathcal{X}$ such that $p = f \cdot \phi$, where $p : Y \times C \rightarrow C$ is the natural projection.

$$\begin{array}{ccc} Y \times C & \xrightarrow{\phi} & \mathcal{X} \\ p \downarrow & & \downarrow f \\ C & \xlongequal{\quad} & C \end{array}$$

Let $\text{Hom}(C, \mathcal{X})$ be a scheme consisting of k -morphisms from C to $\mathcal{X}$. Then, we have the natural morphism of schemes $\alpha : \text{Hom}(C, \mathcal{X}) \rightarrow \text{Hom}(C, C)$ defined by $\alpha(s) = f \cdot s$ for $s \in \text{Hom}(C, \mathcal{X})$. We set $\text{Sec}(C, \mathcal{X}) = \alpha^{-1}(\text{id}_C)$.

By the definition of Σ_A , Σ_A is a bounded family. So we can find finite number of connected components $\text{Sec}_1, \dots, \text{Sec}_l$ of $\text{Sec}(C, \mathcal{X})$ such that, for all $P \in S_A$, there is $s \in \coprod_{i=1}^l \text{Sec}_i$ with $s(C) = \Delta_P$. Further, since every connected component of $\text{Hom}(C, \mathcal{X})$ is quasi-projective, Sec_i 's are also quasi-projective.

On the other hand, since S_A is dense in $X(\bar{F})$, there is a Sec_i such that the natural morphism $\text{Sec}_i \times C \rightarrow \mathcal{X}$ is a dominant morphism. Moreover, since $\mathcal{X}$ is irreducible, there is an irreducible component Y of Sec_i such that $Y \times C \rightarrow \mathcal{X}$ still dominates $\mathcal{X}$. Replacing Y by Y_{red} , we may assume that Y is reduced. Furthermore, considering a desingularization $Y' \rightarrow Y$ of Y , we may also assume that Y is smooth. Thus, we have our claim.

Let Y_1 be a smooth compactification of Y . Then, ϕ induces a rational map $\phi_1 : Y_1 \times C \dashrightarrow \mathcal{X}$. Let $\mu : Z \rightarrow Y_1 \times C$ be a minimal succession of blowing-ups such that $\phi_1 \cdot \mu : Z \rightarrow \mathcal{X}$ gives a morphism. Let E be the exceptional set of μ . Since X contains no rational curves, $(p \cdot \mu)(E)$ is a proper closed subset of C . Therefore, we can take a non-empty open set C_0 of C such that ϕ_1 is defined on $Y_1 \times C_0$. Thus, by virtue of Lemma 1.5, for a general point t of C_0 , the Kodaira-Spencer map $\varrho_t : T_{C,t} \rightarrow H^1(\mathcal{X}_t, T_{\mathcal{X}_t})$ is zero. This is a contradiction. Therefore, S_A is not dense in $X(\bar{F})$. $\square$

Remark 1.6. For geometric heights, finiteness does not hold in general. For, as pointing out in Example of [No], we have a non-isotrivial family of projective varieties with ample cotangent bundles, which contains a trivial family of subvarieties. More precisely, we have the following example with properties (1) — (5).

- (1) Let $f : \mathcal{X} \rightarrow C$ be a projective morphism of smooth varieties.
- (2) $\dim \mathcal{X} = 3$, $\dim C = 1$ and $f_* \mathcal{O}_{\mathcal{X}} = \mathcal{O}_C$.
- (3) $\Omega_{\mathcal{X}_t}^1$ is ample for a general $t \in C$.
- (4) The Kodaira-Spencer map $\varrho_t : T_{C,t} \rightarrow H^1(\mathcal{X}_t, T_{\mathcal{X}_t})$ is injective for a general $t \in C$.
- (5) There are a smooth projective curve C' and an embedding $j : C' \times C \rightarrow \mathcal{X}$ with $f \cdot j = p$, where p is the natural projection $p : C' \times C \rightarrow C$.

$$\begin{array}{ccc} C' \times C & \xrightarrow{j} & \mathcal{X} \\ p \downarrow & & \downarrow f \\ C & \xlongequal{\quad} & C \end{array}$$

2. Ample vector bundles.

In this section, we will prepare basic facts of ample vector bundles. Let us begin with Hartshorne's criterion of ample vector bundles on curves.

Lemma 2.1. (cf. [Ha, Theorem 2.4]) *Let C be a smooth projective curve over an algebraically closed field of characteristic zero and E a vector bundle on C with $\deg(E) > 0$. Then, E is ample if and only if every quotient bundle of E has positive degree.*

Proof. For reader's convenience, we will give a sketch of an elementary proof. Clearly, it is sufficient to show that if every quotient bundle of E has positive degree, then E is ample.

First, let us consider a case where E is semistable. Let $\pi : \mathbb{P}(E) \rightarrow C$ be the projective bundle of E and $\mathcal{O}(1)$ the tautological line bundle of E . Then, by [Mi, Theorem 3.1], $\mathcal{O}(r) \otimes \pi^*(\det(E)^{-1})$ is nef, where $r = \operatorname{rk} E$. Thus, $\mathcal{O}(r)$ is ample because $\deg(E) > 0$.

Next, we deal with a general case. We prove it by induction on $\operatorname{rk} E$. By the previous observation, we may assume that E is not semistable. Let E_1 be the maximal destabilizing subbundle of E . Since E_1 is semistable and $\deg(E_1) > 0$, E_1 is ample as before. On the other hand, by hypothesis of induction, E/E_1 is ample. Thus, E is ample. $\square$

Corollary 2.2. *Let C be a smooth projective curve over an algebraically closed field of characteristic zero and R an ample vector bundle on C . Let*

$$(2.2.1) \quad 0 \rightarrow \mathcal{O}_C \rightarrow E \rightarrow R \rightarrow 0$$

be an extension of R by $\mathcal{O}_C$. Then, E is ample if and only if (2.2.1) does not split.

Proof. First, we assume that E is ample. Let $\pi : \mathbb{P}(E) \rightarrow C$ be the projective bundle of E and $\mathcal{O}(1)$ the tautological line bundle. If (2.2.1) splits, then we have a section Δ of π

defined by a splitting $E \rightarrow \mathcal{O}_C$. Clearly, $(\mathcal{O}(1) \cdot \Delta) = 0$. This is a contradiction because $\mathcal{O}(1)$ is ample. Hence, (2.2.1) doesn't split.

Next we assume that (2.2.1) doesn't split. Let Q be a quotient bundle of E . Let $\alpha : \mathcal{O}_C \rightarrow E \rightarrow Q$ be the composition homomorphism. If α is zero, we have a surjective homomorphism $R \rightarrow Q$. Thus, $\deg(Q) > 0$. So we may assume that $\alpha \neq 0$. Let L be the saturation of $\alpha(\mathcal{O}_C)$ in Q . Let us consider the following homomorphisms:

$$R \simeq E/\mathcal{O}_C \rightarrow Q/\alpha(\mathcal{O}_C) \rightarrow (Q/\alpha(\mathcal{O}_C))/(\text{torsions}) \simeq Q/L.$$

This shows that there is a surjective homomorphism $R \rightarrow Q/L$. Thus, if $\text{rk } Q \geq 2$, then $\deg(Q) = \deg(Q/L) + \deg(L) > 0$. Moreover, in the case where $\text{rk } Q = 1$, since (2.2.1) doesn't split, α is not an isomorphism. Hence, $\deg(Q) = \deg(L) > 0$. Therefore, E is ample by Lemma 2.1. $\square$

Proposition 2.3. *Let X be a smooth projective variety over an algebraically closed field of characteristic zero and Q an ample vector bundle on X . Let $0 \rightarrow \mathcal{O}_X \rightarrow E \rightarrow Q \rightarrow 0$ be an extension of Q by $\mathcal{O}_X$. Let $\pi : P = \mathbb{P}(E) \rightarrow X$ be the projective bundle of E , $\mathcal{O}_P(1)$ the tautological line bundle of P , and M a line bundle on P . If $0 \rightarrow \mathcal{O}_X \rightarrow E \rightarrow Q \rightarrow 0$ does not split, then, for a sufficiently large n , $\pi(\text{Bs}(\mathcal{O}_P(n) \otimes M))$ is a proper closed subset of X .*

Proof. Let Y be a sub-variety of P defined by $E \rightarrow Q$. Then, $Y \in |\mathcal{O}_P(1)|$. First of all, since $\mathcal{O}_P(1)|_Y$ is ample, if $n \gg 0$, then $H^i(Y, \mathcal{O}_P(n) \otimes M|_Y) = 0$ for $i > 0$ and $\mathcal{O}_P(n) \otimes M|_Y$ is generated by global sections. Let us consider an exact sequence:

$$0 \rightarrow \mathcal{O}_P(n-1) \otimes M \rightarrow \mathcal{O}_P(n) \otimes M \rightarrow \mathcal{O}_P(n) \otimes M|_Y \rightarrow 0.$$

If $n \gg 0$,

$$(2.3.1) \quad H^1(P, \mathcal{O}_P(n-1) \otimes M) \rightarrow H^1(P, \mathcal{O}_P(n) \otimes M)$$

is surjective. So, $h^1(P, \mathcal{O}_P(n) \otimes M)$ is constant for $n \gg 0$. Therefore, (2.3.1) is bijective for $n \gg 0$. Hence, we get

$$H^0(P, \mathcal{O}_P(n) \otimes M) \rightarrow H^0(Y, \mathcal{O}_P(n) \otimes M|_Y)$$

is surjective for $n \gg 0$. It follows that, if we set $B_n = \text{Bs}(\mathcal{O}_P(n) \otimes M)$, then $B_n \cap Y = \emptyset$ for $n \gg 0$.

Let H be an ample line bundle on X . If $\dim X \geq 2$, then $H^1(X, Q^\vee \otimes H^{-m}) = 0$ for sufficiently large m . Thus, choosing a general member T of $|H^m|$, the restriction map $H^1(X, Q^\vee) \rightarrow H^1(T, Q^\vee|_T)$ is injective. This observation shows us that, for a general curve C on X , $H^1(X, Q^\vee) \rightarrow H^1(C, Q^\vee|_C)$ is injective. This means that the exact sequence:

$$0 \rightarrow \mathcal{O}_C \rightarrow E|_C \rightarrow Q|_C \rightarrow 0$$

does not split for the curve C . Hence, by Corollary 2.2, $E|_C$ is ample.

Here we assume that $\pi(B_n) = X$. Then, there is an integral curve C' such that $C' \subset B_n$ and $\pi(C') = C$. Then, since $C' \cap Y = \emptyset$, we have $(\mathcal{O}_P(1) \cdot C') = 0$. This contradicts to ampleness of $E|_C$. $\square$

3. Proofs of Theorem A and Theorem B.

3.1. Proof of Theorem A. Let $(f : \mathcal{X} \rightarrow C, \mathcal{L})$ be a model of (X, L) . Let $\pi : \mathcal{Y} = \mathbb{P}(\Omega_{\mathcal{X}/k}^1) \rightarrow \mathcal{X}$ be the projective bundle and $\mathcal{O}_{\mathcal{Y}}(1)$ the tautological line bundle on $\mathcal{Y}$. We set $Y = \mathcal{Y}_F$, $B'_n = \text{Bs}(\mathcal{O}_Y(n) \otimes \pi_F^*(L^{-1}))$, and $B_n = \pi_F(B'_n)$. By Proposition 1.2, there is a constant D such that, for all $P' \in Y(\bar{F}) \setminus B'_n(\bar{F})$,

$$(3.1.1) \quad h_{\mathcal{O}_Y(n) \otimes \pi_F^*(L^{-1})}(P') \geq D.$$

On the other hand, the canonical exact sequence:

$$0 \rightarrow f^*(\Omega_{C/k}^1) \rightarrow \Omega_{\mathcal{X}/k}^1 \rightarrow \Omega_{\mathcal{X}/C}^1 \rightarrow 0$$

gives rise to an exact sequence on X :

$$(3.1.2) \quad 0 \rightarrow \mathcal{O}_X \rightarrow \Omega_{\mathcal{X}/k}^1 \Big|_X \rightarrow \Omega_{X/F}^1 \rightarrow 0.$$

Since X is non-isotrivial, (3.1.2) does not split. Thus, by virtue of Proposition 2.3, if $n \gg 0$, then $B_n \otimes \bar{F}$ is a proper closed subset of $X_{\bar{F}}$. Therefore, so is B_n .

Let $P \in X(\bar{F}) \setminus B_n(\bar{F})$, Δ_P the corresponding integral curve on $\mathcal{X}$, $\tilde{\Delta}_P$ the normalization of Δ_P and $\iota : \tilde{\Delta}_P \rightarrow \mathcal{X}$ the natural morphism. Moreover, let $\phi : \mathbb{P}(\iota^*(\Omega_{\mathcal{X}/k}^1)) \rightarrow \mathcal{Y}$ be the induced morphism.

$$\begin{array}{ccc} \mathbb{P}(\iota^*(\Omega_{\mathcal{X}/k}^1)) & \xrightarrow{\phi} & \mathcal{Y} = \mathbb{P}(\Omega_{\mathcal{X}/k}^1) \\ \downarrow & & \downarrow \pi \\ \tilde{\Delta}_P & \xrightarrow{\iota} & \mathcal{X} \end{array}$$

Let Q be the minimal quotient line bundle of $\iota^*(\Omega_{\mathcal{X}/k}^1)$ and T the corresponding section of $\mathbb{P}(\iota^*(\Omega_{\mathcal{X}/k}^1))$. Let $\Delta' = \phi(T)$ and P' the corresponding $\bar{F}$ -value point of Y . Then, $P' \in Y(\bar{F}) \setminus B'_n(\bar{F})$. Thus, by (3.1.1),

$$(3.1.3) \quad D \leq h_{\mathcal{O}_Y(n) \otimes \pi_F^*(L^{-1})}(P') = n \frac{\deg(Q)}{[F(P) : F]} - h_L(P).$$

On the other hand, since there is a non-trivial homomorphism $\iota^*(\Omega_{\mathcal{X}/k}^1) \rightarrow \Omega_{\tilde{\Delta}_P/k}^1$, we have $\deg Q \leq \deg(\Omega_{\tilde{\Delta}_P/k}^1)$. Thus, by (3.1.3), we get $h_L(P) \leq n \cdot d(P) - D$. $\square$

3.2. Proof of Theorem B. By Theorem A and Theorem 1.3, it is sufficient to see the following lemma.

Lemma 3.2.1. *Let X be a smooth projective variety over a field K with the ample cotangent bundle and C a 1-dimensional subvariety of X . Then, the genus of C is greater than one.*

Proof. Let $\tilde{C}$ be the normalization of C and $\phi : \tilde{C} \rightarrow X$ the induced morphism. Then, we have a non-trivial homomorphism $\phi^*(\Omega_{X/K}^1) \rightarrow \Omega_{\tilde{C}/K}^1$. Here, $\phi^*(\Omega_{X/K}^1)$ is ample. Therefore, $\deg(\Omega_{\tilde{C}/K}^1) > 0$. $\square$

REFERENCES

- [Ha] R. Hartshorne, *Ample vector bundles on curves*, Nagoya Math. J. **43** (1971), 73–89.
- [La] S. Lang (Ed.), *Encyclopaedia of Mathematical Sciences, Vol. 60, Number Theory III.*, Springer-Verlag, 1991.
- [Mi] Y. Miyaoka, *The Chern classes and Kodaira dimension of a minimal variety*, Advanced Studies in Pure Mathematics 10, 1987, Algebraic Geometry, Sendai, 1985, Kinokuniya, 449–476.
- [Mo] A. Moriwaki, *Height inequality of non-isotrivial curves over function fields*, to appear in J. Algebraic Geometry.
- [No] J. Noguchi, *A higher dimensional analogue of Mordell’s conjecture over function fields*, Math. Ann. **258** (1981), 207–212.
- [Vo] P. Vojta, *On algebraic points on curves*, Comp. Math. **78** (1991), 29–36.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KYOTO UNIVERSITY, KYOTO, 606-01,
JAPAN

Current address: Department of Mathematics, University of California, Los Angeles, 405 Hilgard
Avenue, Los Angeles, California 90024, USA

E-mail address: moriwaki@math.ucla.edu